

# Some results *à l'Abel* obtained by use of techniques *à la Hopf*

Christian Costermans

Hoang Ngoc Minh

University Lille II, 1, Place Déliot, 59024 Lille, [ccostermans,hoang]@univ-lille2.fr

## Abstract

In this work, we obtain some results *à l'Abel* dealing with noncommutative generating series of polylogarithms and multiple harmonic sums, by using techniques *à la Hopf*. In particular, this enables to explicit generalized Euler constants associated to divergent polyzêtas and to extract the constant part of (commutative and noncommutative) generating series of all polyzêtas.

**Keywords:** asymptotic expansion, generating series, multiple harmonic sums, polylogarithms, polyzêtas

## 1 Introduction

Let us consider the alphabet  $Y = \{y_i\}_{i \in \mathbb{N}_+}$ . To each word  $w = y_{s_1} \dots y_{s_r}$  of the monoid  $Y^*$ , we associate the multiple harmonic sum  $H_w(N)$  and the polylogarithm  $\text{Li}_w(z)$

$$H_w(N) = \sum_{N \geq n_1 > \dots > n_r > 0} \frac{1}{n_1^{s_1} \dots n_r^{s_r}}, \quad \text{Li}_w(z) = \sum_{n_1 > \dots > n_r > 0} \frac{z^{n_1}}{n_1^{s_1} \dots n_r^{s_r}}. \quad (1)$$

For  $0 \leq N < r$ ,  $H_w(N) = 0$  and for the empty word  $\epsilon$ , we put  $H_\epsilon(N) = 1$ , for any  $N \geq 0$ . For  $w \in Y^* \setminus y_1 Y^*$ , the limits  $\lim_{z \rightarrow 1} \text{Li}_w(z)$  and  $\lim_{N \rightarrow \infty} H_w(N)$  exist and, by an Abel theorem, are equal to the convergent polyzêta  $\zeta(w)$

$$\zeta(w) = \sum_{n_1 > \dots > n_r > 0} \frac{1}{n_1^{s_1} \dots n_r^{s_r}}, \quad s_1 > 1. \quad (2)$$

In other cases, *i.e.* for  $w = y_s w'$ , the associated polylogarithm  $\text{Li}_w$  and polyzêta  $\zeta(w)$  can be considered respectively as a polylogarithmic generating series and as Dirichlet series

$$\text{Li}_{y_s w'}(z) = \sum_{N > 0} \frac{p_N}{N^s} z^N \quad \text{and} \quad \zeta(y_s w') = \sum_{N > 0} \frac{p_N}{N^s} \quad (3)$$

with  $p_N = H_{w'}(N - 1)$ . Both series can be obtained from the following generating series

$$P_{w'}(z) = \sum_{N=0}^{\infty} H_{w'}(N) z^N = \sum_{N \geq 0} p_{N+1} z^N, \quad (4)$$

respectively by the polylogarithmic transform and by the Mellin transform [?]

$$\text{Li}_{y_s w'}(z) = \int_0^{\infty} \frac{P_{w'}(ze^{-u})}{\Gamma(s)} \frac{du}{u^{1-s}} \quad \text{and} \quad \zeta(y_s w') = \int_0^{\infty} \frac{P_{w'}(e^{-u})}{\Gamma(s)} \frac{du}{u^{1-s}}. \quad (5)$$

Therefore, the knowledge of the singular expansion of the ordinary generating series  $P_{w'}(z) = (1 - z)^{-1} \text{Li}_{w'}(z)$  in the scale  $\{(1 - z)^a \log^b(1 - z)\}_{a \in \mathbb{Z}, b \in \mathbb{N}}$  enables to get, on one hand the asymptotic behaviour, as  $N \rightarrow \infty$ , of its Taylor coefficients  $H_{w'}(N)$  in the scale  $\{N^\alpha \log^\beta N\}_{\alpha \in \mathbb{Z}, \beta \in \mathbb{N}}$ . Hence, to deduce the behavior of  $H_w(N) = \sum_{i=1}^N H_{w'}(i - 1)/i^s$ . This gives on the other hand, through tauberian theorem, the singular expansion of Dirichlet series  $\zeta(y_s w')$  considered then as a function of the complex variable  $s$ .

Both studies lead to apply another Abel theorem dealing with Dirichlet series [?]. Indeed, let us consider the partial sum  $S_N$  of coefficients of the ordinary generating series  $P_{w'}(z)$ ,

$$S_N = \sum_{i=1}^N p_{i+1} = \sum_{i=1}^N H_{w'}(i). \quad (6)$$

If  $S_N$  admits a singular expansion of the following type

$$S_N = \sum_{j=1}^k B_j N^{\sigma_j} \log^{\alpha_j} N + O(N^\beta), \quad (7)$$

where, for all  $j = 1, \dots, k$ ,  $B_j$  is an arbitrary complex number,  $\sigma_j, \alpha_j$  are arbitrary integers, and  $\beta$  is an integer such that  $\beta > \sigma_k$ , then the Dirichlet series  $\zeta(y_s w')$  is convergent for  $s > 1$  and even regular except in  $\sigma_1, \dots, \sigma_k$  which are its logarithmic singularities.

In order to adapt automatically these Abel techniques to polylogarithms  $\{\text{Li}_w\}_{w \in Y^*}$  and to multiple harmonic sums  $\{H_w\}_{w \in Y^*}$ , we consider the *noncommutative generating series*

$$\Lambda(z) = \sum_{w \in Y^*} \text{Li}_w(z) w \quad \text{and} \quad H(N) = \sum_{w \in Y^*} H_w(N) w. \quad (8)$$

Through algebraic combinatoric [?] and elements of topology of formal series in noncommutative variables [?], we show in Section ?? the existence of formal series over  $Y$ ,  $Z_1$  and  $Z_2$  in non commutative variables with constant coefficients, such that

$$\lim_{z \rightarrow 1} \exp \left[ y_1 \log \frac{1}{1 - z} \right] \Lambda(z) = Z_1 \quad \text{and} \quad \lim_{N \rightarrow \infty} \exp \left[ \sum_{k \geq 1} H_{y_k}(N) \frac{(-y_1)^k}{k} \right] H(N) = Z_2. \quad (9)$$

Moreover, we have  $Z_1 = Z_2$ , both standing for the noncommutative generating series of all convergent polyzêtas  $\{\zeta(w)\}_{w \in Y^* \setminus y_1 Y^*}$ , (as shown by the factorized form). This enables, in particular, to explicit generalized Euler constants associated to divergent polyzêtas  $\{\zeta(w)\}_{w \in y_1 Y^*}$  and to extract the constant part of generating series (commutative and noncommutative) of all polyzêtas.

Techniques presented in this paper can be applied to other fields, like polysystems occurring in physical problems, and enable to make the calculations easier. To illustrate this, we present in appendix some results to compute, thanks to such techniques, the solution of a linear differential system, with three singularities, that can be supposed to be  $\{0, 1, \infty\}$ , after an homographic transformation.

## 2 Polylogarithm and harmonic sum

### 2.1 Algebraic properties

#### 2.1.1 Symmetric functions and harmonic sums

Let  $\{t_i\}_{i \in \mathbb{N}_+}$  be an infinite set of variables. The elementary symmetric functions  $\lambda_k$  and the sums of powers  $\psi_k$  are defined by

$$\lambda_k(\underline{t}) = \sum_{n_1 > \dots > n_k > 0} t_{n_1} \dots t_{n_k} \quad \text{and} \quad \psi_k(\underline{t}) = \sum_{n > 0} t_n^k. \quad (10)$$

They are respectively coefficients of the following generating functions

$$\lambda(\underline{t}|z) = \sum_{k > 0} \lambda_k(\underline{t}) z^k = \prod_{i \geq 1} (1 + t_i z) \quad \text{and} \quad \psi(\underline{t}|z) = \sum_{k > 0} \psi_k(\underline{t}) z^{k-1} = \sum_{i \geq 1} \frac{t_i}{1 - t_i z}. \quad (11)$$

These generating functions satisfy a Newton identity

$$d/dz \log \lambda(\underline{t}|z) = \psi(\underline{t}| -z). \quad (12)$$

The fundamental theorem from symmetric functions theory asserts that the  $\{\lambda_k\}_{k \geq 0}$  are linearly independent, and remarkable identities give (putting  $\lambda_0 = 1$ ) :

$$k! \lambda_k = (-1)^k \sum_{\substack{s_1, \dots, s_k > 0 \\ s_1 + \dots + k s_k = k}} \binom{k}{s_1, \dots, s_k} \left(-\frac{\psi_1}{1}\right)^{s_1} \dots \left(-\frac{\psi_k}{k}\right)^{s_k} \quad (13)$$

Let  $w = y_{s_1} \dots y_{s_r} \in Y^*$ . The quasi-symmetric function  $F_w$ , of depth  $r = |w|$  and of degree (or weight)  $s_1 + \dots + s_r$ , is defined by

$$F_w(\underline{t}) = \sum_{n_1 > \dots > n_r > 0} t_{n_1}^{s_1} \dots t_{n_r}^{s_r}. \quad (14)$$

In particular,  $F_{y_1^k} = \lambda_k$  and  $F_{y_k} = \psi_k$ . As a consequence, the functions  $\{F_{y_1^k}\}_{k \geq 0}$  are linearly independent and integrating differential equation (??) shows that functions  $F_{y_1^k}$  and  $F_{y_k}$  are linked by the formula

$$\sum_{k \geq 0} F_{y_1^k} z^k = \exp \left[ - \sum_{k \geq 1} F_{y_k} \frac{(-z)^k}{k} \right]. \tag{15}$$

Remarkable identity (??) can be then seen as

$$k! y_1^k = (-1)^k \sum_{\substack{s_1, \dots, s_k > 0 \\ s_1 + \dots + k s_k = k}} \binom{k}{s_1, \dots, s_k} \frac{(-y_1)^{\sqcup s_1}}{1^{s_1}} \sqcup \dots \sqcup \frac{(-y_k)^{\sqcup s_k}}{k^{s_k}} \tag{16}$$

Every  $H_w(N)$  can be obtained by specializing variables  $\{t_i\}_{N \geq i \geq 1}$  at  $t_i = 1/i$  and, for  $i > N, t_i = 0$  in the quasi-symmetric function  $F_w$  [?]. In the same way, when  $w \in Y^* \setminus y_1 Y^*$ , the convergent polyzêta  $\zeta(w)$  can be obtained by specializing variables  $\{t_i\}_{i \geq 1}$  at  $t_i = 1/i$  in  $F_w$  [?]. The notation  $F_w$  is extended by linearity to all polynomials over  $Y$ .

If  $u$  (resp.  $v$ ) is a word in  $Y^*$ , of length  $r$  and of weight  $p$  (resp. of length  $s$  and of weight  $q$ ),  $F_{u \sqcup v}$  is a quasi-symmetric function of depth  $r + s$  and of weight  $p + q$ , and we have  $F_{u \sqcup v} = F_u F_v$ . In consequence,  $H_{u \sqcup v} = H_u H_v$  [?]. In the same way, when  $u, v \in Y^* \setminus y_1 Y^*$ , we also have  $\zeta(u \sqcup v) = \zeta(u) \zeta(v)$  [?].

Let us consider the noncommutative generating series  $H(N)$  of  $\{H_w(N)\}_{w \in Y^*}$  [?],

$$H(N) = \sum_{w \in Y^*} H_w(N) w = \prod_1^{l=N} \left( 1 + \sum_{i > 0} \frac{y_i}{l^i} \right). \tag{17}$$

since it verifies the difference equation

$$H(N) = \left( 1 + \sum_{i > 0} \frac{y_i}{N^i} \right) H(N - 1) \quad \text{with} \quad H(0) = 1. \tag{18}$$

### 2.1.2 Polylogarithms and polyzêtas

Let us denote by  $\mathcal{C}$  the algebra  $\mathbb{C}[z, 1/z, 1/(1 - z)]$  of polynomial functions in  $z, 1/z$  and  $1/(1 - z)$ . We define two differential forms  $\omega_0(z) = dz/z$  and  $\omega_1(z) = dz/(1 - z)$ .

Let  $w = x_0^{s_1-1} x_1 \dots x_0^{s_r-1} x_1 \in X^* x_1$ . One can check that the polylogarithm  $\text{Li}_w$  is also the value of the iterated integral over  $\omega_0, \omega_1$  and along the integration path  $0 \rightsquigarrow z$  :

$$\text{Li}_w(z) = \int_{0 \rightsquigarrow z} \omega_0^{s_1-1} \omega_1 \dots \omega_0^{s_r-1} \omega_1. \tag{19}$$

This provides an analytic continuation of the  $\text{Li}_w$  over the universal covering  $\widetilde{\mathbb{C} - \{0, 1\}}$  of  $\mathbb{C}$  without points 0 and 1. We extend the definition of polylogarithms over  $X^*$  putting

$$\text{Li}_{x_0^k}(z) = \log^k z/k!, \quad \text{for} \quad k \in \mathbb{N}. \tag{20}$$

Let  $\text{LI}_{\mathcal{C}} = (\mathcal{C}\{\text{Li}_w\}_{w \in X^*}, \cdot)$  be the smallest  $\mathcal{C}$ -algebra containing  $\mathcal{C}$  and stable by differentiation and integration over  $\omega_0, \omega_1$ . It can be identified with the  $\mathcal{C}$ -module generated by polylogarithms. Thus, the polylogarithms are  $\mathcal{C}$ -linearly independent [?]. Hence,  $(\mathcal{C}\{\text{Li}_w\}_{w \in X^*}, \cdot)$  is identified with the polynomial algebra  $(\mathcal{C}\{P_l\}_{l \in \mathcal{L}ynX}, \cdot)$  [?]. The noncommutative generating series  $L = \sum_{w \in X^*} \text{Li}_w$   $w$  satisfies Drinfel'd differential equation [?, ?]

$$dL = (x_0\omega_0 + x_1\omega_1)L \quad \text{with the condition } L(\varepsilon) = e^{x_0 \log \varepsilon} + O(\sqrt{\varepsilon}) \quad \text{for } \varepsilon \rightarrow 0^+. \quad (21)$$

This enables to prove that  $L$  is the exponential of a Lie series [?]. So, applying a Ree theorem, it verifies Friedrichs criterion [?], *i.e.*  $\text{Li}_{u \sqcup v} = \text{Li}_u \text{Li}_v$ , for  $u, v \in X^*$ . In particular, when  $u, v \in x_0X^*x_1$ , we also have  $\zeta(u \sqcup v) = \zeta(u) \zeta(v)$ . From the factorization of monoid by Lyndon words, we obtain the factorization of the series  $L$  [?]:

$$L(z) = e^{x_1 \log \frac{1}{1-z}} \left[ \prod_{l \in \mathcal{L}ynX \setminus \{x_0, x_1\}} e^{\text{Li}_{S_l}(z)[l]} \right] e^{x_0 \log z}. \quad (22)$$

For all  $l \in \mathcal{L}ynX \setminus \{x_0, x_1\}$ , we have  $S_l \in x_0X^*x_1$ . So, let us put [?]

$$L_{\text{reg}} = \prod_{l \in \mathcal{L}ynX \setminus \{x_0, x_1\}} e^{\text{Li}_{S_l}[l]} \quad \text{and} \quad Z = L_{\text{reg}}(1). \quad (23)$$

Let  $\sigma$  be the monoid endomorphism verifying  $\sigma(x_0) = -x_1, \sigma(x_1) = -x_0$ , we also get [?]

$$L(z) = \sigma[L(1-z)]Z = e^{x_0 \log z} \sigma[L_{\text{reg}}(1-z)]e^{-x_1 \log(1-z)}Z. \quad (24)$$

In consequence, from (??) and (??), we get respectively

$$L(z) \underset{z \rightarrow 0}{\sim} \exp(x_0 \log z) \quad \text{and} \quad L(z) \underset{z \rightarrow 1}{\sim} \exp\left(x_1 \log \frac{1}{1-z}\right)Z. \quad (25)$$

Let  $\pi_Y : \text{LI}_{\mathcal{C}}\langle\langle X \rangle\rangle \rightarrow \text{LI}_{\mathcal{C}}\langle\langle Y \rangle\rangle$  a projector s.t., for  $f \in \text{LI}_{\mathcal{C}}, w \in X^*, \pi_Y(f w x_0) = 0$ . Then

$$\Lambda(z) = \pi_Y L(z) \underset{z \rightarrow 1}{\sim} \exp\left(y_1 \log \frac{1}{1-z}\right) \pi_Y Z. \quad (26)$$

**Definition 1** ([?]). Let  $\zeta_{\sqcup} : (\mathcal{C}\langle\langle X \rangle\rangle, \sqcup) \rightarrow (\mathbb{C}, \cdot)$  be the algebra morphism (*i.e.* for  $u, v \in X^*, \zeta_{\sqcup}(u \sqcup v) = \zeta_{\sqcup}(u)\zeta_{\sqcup}(v)$ ) verifying for all convergent word  $w \in x_0X^*x_1, \zeta_{\sqcup}(w) = \zeta(w)$ , and such that  $\zeta_{\sqcup}(x_0) = \zeta_{\sqcup}(x_1) = 0$ .

Then, the noncommutative generating series  $Z_{\sqcup} = \sum_{w \in X^*} \zeta_{\sqcup}(w) w$  verifies  $Z_{\sqcup} = Z$  [?]. In consequence,  $Z_{\sqcup}$  is the unique Lie exponential verifying  $\langle Z_{\sqcup} | x_0 \rangle = \langle Z_{\sqcup} | x_1 \rangle = 0$

and  $\langle Z_{\sqcup} | w \rangle = \zeta(w)$ , for any  $w \in x_0 X^* x_1$ . Its logarithm is given by  $\log Z_{\sqcup} = \sum_{w \in X^*} \zeta_{\sqcup}(w) \pi_1(w)$ , where  $\pi_1(w)$  is the Lie polynomial [?]

$$\pi_1(w) = \sum_{k \geq 1} \frac{(-1)^{k-1}}{k} \sum_{u_1, \dots, u_k \in X^* \setminus \{\epsilon\}} \langle w | u_1 \sqcup \dots \sqcup u_k \rangle u_1 \cdots u_k. \quad (27)$$

The series  $Z$  shall be understood then as Drinfel'd associator  $\Phi_{KZ}$  [?, ?] verifying duality, pentagonal and hexagonal relations. We also can obtain the expression of this associator given by L e and Murakami [?] thanks to the following expansion [?]

$$Z = \sum_{k \geq 0} \sum_{l_1, \dots, l_k \geq 0} \zeta_{\sqcup}(x_1 x_0^{l_1} \circ \dots \circ x_1 x_0^{l_k}) \prod_{i=1}^k \text{ad}_{x_0}^{l_i} x_1, \quad (28)$$

where  $\text{ad}_{x_0}^l x_1$  stands for the iterated Lie bracket  $[x_0, [\dots, [x_0, x_1] \dots]]$ , for  $l > 0$  and  $\text{ad}_{x_0}^0 x_1 = x_1$ , the operation  $\circ$  being defined as  $x_1 x_0^l \circ P = x_1(x_0^l \sqcup P)$ , for any  $P \in \mathbb{C}\langle X \rangle$ .

### 2.1.3 Ordinary generating series of harmonic sums

The functions  $\{Li_w\}_{w \in X^*}$  are  $\mathbb{C}$ -linearly independent. Thus, the functions  $\{P_w\}_{w \in Y^*}$  are also  $\mathbb{C}$ -linearly independent and then, the functions  $\{H_w\}_{w \in Y^*}$  are also  $\mathbb{C}$ -linearly independent. So,

**Proposition 1** ([?]). *Extended by linearity, the application  $P : u \mapsto P_u$  is an isomorphism from polynomial algebra  $(\mathbb{C}\langle Y \rangle, \sqcup)$  over Hadamard algebra  $(\mathbb{C}\{P_w\}_{w \in Y^*}, \odot)$ . Moreover, the application  $H : u \mapsto H_u = \{H_u(N)\}_{N \geq 0}$  is an isomorphism from  $(\mathbb{C}\langle Y \rangle, \sqcup)$  over the algebra  $(\mathbb{C}\{H_w\}_{w \in Y^*}, \cdot)$ .*

*Proof*: Indeed, on the first hand,  $\ker P = \{0\}$  and  $\ker H = \{0\}$ , and on the other hand,  $P$  is a morphism for Hadamard product (it inherits of  $H$  for the harmonic product) :

$$\begin{aligned} P_u(z) \odot P_v(z) &= \sum_{N \geq 0} H_u(N) H_v(N) z^N \\ &= \sum_{N \geq 0} H_{u \sqcup v}(N) z^N \\ &= P_{u \sqcup v}(z). \end{aligned}$$

Studying the equivalence between action of  $\{(1 - z)^l\}_{l \in \mathbb{Z}}$  over  $\{P_w(z)\}_{w \in Y^*}$  and this of  $\{N^k\}_{k \in \mathbb{Z}}$  over  $\{H_w(N)\}_{w \in Y^*}$  [?], we have

**Theorem 1.** *The Hadamard  $\mathbb{C}$ -algebra of  $\{P_w\}_{w \in Y^*}$  can be identified with this of  $\{P_l\}_{l \in \mathcal{L}_{yn} Y}$ . Identically, the algebra of harmonic sums  $\{H_w\}_{w \in Y^*}$  with polynomial coefficients can be identified with this of  $\{H_l\}_{l \in \mathcal{L}_{yn} Y}$ .*

As for polylogarithms, we extend the definition of  $P_w$  putting  $P_w(z) = (1-z)^{-1} \text{Li}_w(z)$ , for any  $w \in X^*$ . The noncommutative generating series of  $\{P_w\}_{w \in X^*}$  is defined by

$$P(z) = \sum_{w \in X^*} P_w(z) w = \frac{L(z)}{1-z}. \quad (29)$$

In consequence, by (??), we have

$$P(z) = e^{-(x_1+1) \log(1-z)} L_{\text{reg}}(z) e^{x_0 \log z}. \quad (30)$$

**Lemma 1.** *Let  $\text{Mono}(z) = e^{-(x_1+1) \log(1-z)}$ . Then*

$$\text{Mono} = \sum_{k \geq 0} P_{y_1^k} y_1^k \quad \text{and} \quad \text{Mono}^{-1} = \sum_{k \geq 0} P_{y_1^k} (-y_1)^k.$$

Since the coefficient of  $z^N$  in the Taylor expansion of  $P_{y_1^k}$  is  $H_{y_1^k}(N)$  then

**Lemma 2.** *Let  $\text{Const} = \sum_{k \geq 0} H_{y_1^k} y_1^k$ . Then*

$$\text{Const} = \exp \left[ - \sum_{k \geq 1} H_{y_1^k} \frac{(-y_1)^k}{k} \right] \quad \text{and} \quad \text{Const}^{-1} = \exp \left[ \sum_{k \geq 1} H_{y_1^k} \frac{(-y_1)^k}{k} \right].$$

*Proof:* This is a consequence of Formula (??).

**Proposition 2.** *For  $k > 0$ ,  $H_{y_1^k}$  is polynomial in  $\{H_{y_r}\}_{1 \leq r \leq k}$  (which are algebraically independent), and*

$$H_{y_1^k} = \sum_{\substack{s_1, \dots, s_k > 0 \\ s_1 + \dots + k s_k = k}} \frac{(-1)^k}{s_1! \dots s_k!} \left( -\frac{H_{y_1}}{1} \right)^{s_1} \dots \left( -\frac{H_{y_k}}{k} \right)^{s_k}.$$

*Proof:* From Identity (??), and applying the isomorphism  $H$  on the set of Lyndon words  $\{y_r\}_{1 \leq r \leq k}$ , we get the expected result.

**Example 1.**  $H_{y_1^2} = (H_{y_1}^2 - H_{y_2})/2$ ,  $H_{y_1^3} = (H_{y_1}^3 - 3H_{y_2}H_{y_1} + 2H_{y_3})/6$ .

**Proposition 3** ([?]). *Let  $\sigma$  be the morphism verifying  $\sigma(x_0) = -x_1$ ,  $\sigma(x_1) = -x_0$ .*

$$P(z) = e^{x_0 \log z} \left[ \prod_{l \in \mathcal{Lyn} X, \{x_0, x_1\}}^{\searrow} e^{\text{Li}_{S_l}(1-z)\sigma([l])} \right] \text{Mono}(z) Z,$$

*Proof:* On the first hand, from (??) and on the other hand, from (??), we get  $P(z) = e^{x_0 \log z} \sigma[L_{\text{reg}}(1-z)] e^{-(x_1+1) \log(1-z)} Z$ . Using the expressions of  $L_{\text{reg}}(1-z)$  and of  $\text{Mono}(z)$ , we get the expected results.

## 2.2 Asymptotic expansion

### 2.2.1 Results à l'Abel for generating series

**Proposition 4.**  $P(z) \underset{z \rightarrow 0}{\sim} e^{x_0 \log z}$  and  $P(z) \underset{z \rightarrow 1}{\sim} \text{Mono}(z)Z$ .

*Proof:* From  $P(z) = e^{-(x_1+1)\log(1-z)}L_{\text{reg}}(z)e^{x_0 \log z}$ , we can deduce the behaviour of  $P(z)$  around 0. From Formula (??), we get the behaviour of  $P(z)$  around 1.

**Corollary 1.** Let  $\Pi(z) = \pi_Y P(z) = \sum_{w \in Y^*} P_w(z)$  w. Then  $\Pi(z) \underset{z \rightarrow 1}{\sim} \text{Mono}(z)\pi_Y Z$ .

From this, we extract, taking care of Lemma ??, Taylor coefficients of  $P_w$ , and we get

**Corollary 2.**  $H(N) \underset{N \rightarrow \infty}{\sim} \text{Const}(N)\pi_Y Z$ .

**Theorem 2.**

$$\lim_{z \rightarrow 1} \exp\left(y_1 \log \frac{1}{1-z}\right) \Lambda(z) = \lim_{N \rightarrow \infty} \exp\left(\sum_{k \geq 1} H_{y_k}(N) \frac{(-y_1)^k}{k}\right) H(N) = \pi_Y Z.$$

*Proof:* This is a consequence of Formula (??), of Lemma ?? and of Corollary ??. From Proposition ??, we deduce

$$P(z) = e^{x_0 \log z} \left[ \prod_{\substack{l \in \mathcal{L}_{Y^* X}, \\ l \neq x_0, x_1}} z \left( \sum_{k \geq 0} P_{S_l^{\perp k}}(1-z) \frac{(\sigma([l])^k)}{k!} \right) \right] \text{Mono}(z)Z. \quad (31)$$

Hence, the knowledge of Taylor expansion around 0 of  $\{P_w(1-z)\}_{w \in X^*}$  gives

**Theorem 3** ([?]). For all  $g \in \mathcal{C}\{P_w\}_{w \in Y^*}$ , there exist algorithmically computable coefficients  $c_j \in \mathbb{C}$ ,  $\alpha_j \in \mathbb{Z}$  and  $\beta_j \in \mathbb{N}$  such that

$$g(z) \sim \sum_{j=0}^{+\infty} c_j (1-z)^{\alpha_j} \log^{\beta_j}(1-z) \quad \text{for } z \rightarrow 1.$$

In consequence, there exist algorithmically computable coefficients  $b_i \in \mathbb{C}$ ,  $\eta_i \in \mathbb{Z}$  and  $\kappa_i \in \mathbb{N}$  such that

$$[z^n]g(z) \sim \sum_{i=0}^{+\infty} b_i n^{\eta_i} \log^{\kappa_i}(n) \quad \text{for } n \rightarrow \infty.$$

**Corollary 3** ([?]). Let  $\mathcal{Z}$  the  $\mathbb{Q}$ -algebra generated by convergent polyzêtas and  $\mathcal{Z}'$  the  $\mathbb{Q}[\gamma]$ -algebra generated by  $\mathcal{Z}$ . Then there exist algorithmically computable coefficients  $c_j \in \mathcal{Z}$ ,  $\alpha_j \in \mathbb{Z}$  and  $\beta_j \in \mathbb{N}$  such that

$$\forall w \in Y^*, P_w(z) \sim \sum_{j=0}^{+\infty} c_j (1-z)^{\alpha_j} \log^{\beta_j}(1-z) \quad \text{for } z \rightarrow 1.$$

In consequence, there exist algorithmically computable coefficients  $b_i \in \mathcal{Z}'$ ,  $\kappa_i \in \mathbb{N}$  and  $\eta_i \in \mathbb{Z}$  such that

$$\forall w \in Y^*, H_w(N) \sim \sum_{i=0}^{+\infty} b_i N^{\eta_i} \log^{\kappa_i}(N) \quad \text{for } N \rightarrow +\infty.$$

### 2.2.2 Generalized Euler constants associated to divergent polyzêtas

**Definition 2.** Let  $\zeta_{\sqcup} : (\mathbb{C}\langle Y \rangle, \sqcup) \rightarrow (\mathbb{C}, \cdot)$  the algebra morphism (i.e. for all convergent word  $u, v \in Y^*$ ,  $\zeta_{\sqcup}(u \sqcup v) = \zeta_{\sqcup}(u)\zeta_{\sqcup}(v)$ ) verifying for  $w \in Y^* \setminus y_1 Y^*$ ,  $\zeta_{\sqcup}(w) = \zeta(w)$  and such that  $\zeta_{\sqcup}(y_1) = \gamma$ .

**Proposition 5.**

$$\zeta_{\sqcup}(y_1^k) = \sum_{\substack{s_1, \dots, s_k > 0 \\ s_1 + \dots + s_k = k}} \frac{(-1)^k}{s_1! \dots s_k!} (-\gamma)^{s_1} \left(-\frac{\zeta(2)}{2}\right)^{s_2} \dots \left(-\frac{\zeta(k)}{k}\right)^{s_k}.$$

*Proof:* By (??) and applying the (surjective) morphism  $\zeta_{\sqcup}$ , we get the expected result. In consequence,

**Theorem 4** ([?]). For  $k > 0$ , the constant  $\zeta_{\sqcup}(y_1^k)$  associated to divergent polyzêta  $\zeta(y_1^k)$  is a polynomial of degree  $k$  in  $\gamma$  with coefficients in  $\mathbb{Q}[\zeta(2), \zeta(2i+1)]_{0 < i \leq (k-1)/2}$ . Moreover, for  $l = 0, \dots, k$ , the coefficient of  $\gamma^l$  is of weight  $k - l$ .

**Example 2.**  $\zeta_{\sqcup}(y_1^2) = [\gamma^2 - \zeta(2)]/2$ ,  $\zeta_{\sqcup}(y_1^3) = [\gamma^3 - 3\zeta(2)\gamma + 2\zeta(3)]/6$  and  $\zeta_{\sqcup}(y_1^4) = [80\zeta(3)\gamma - 60\zeta(2)\gamma^2 + 6\zeta(2)^2 + 10\gamma^4]/240$ .

Let us consider the (exponential) partial Bell polynomials in the variables  $\{t_l\}_{l \geq 1}$ ,  $b_{n,k}(t_1, \dots, t_{n-k+1})$ , defined by the exponential generating series :

$$\sum_{n=0}^{\infty} \sum_{k=0}^n b_{n,k}(t_1, \dots, t_{n-k+1}) \frac{v^n u^k}{n!} = \exp\left(u \sum_{l=1}^{\infty} t_l \frac{v^l}{l!}\right). \quad (32)$$

In particular, we have

**Lemma 3.** Let  $t_m = (-1)^m (m-1)! \zeta_{\sqcup}(m)$ , for  $m \geq 1$ . Then

$$\exp\left[\sum_{k \geq 1} \zeta_{\sqcup}(k) \frac{(-y_1)^k}{k}\right] = 1 + \sum_{n \geq 1} \left[\sum_{k=1}^n b_{n,k}(\gamma, \zeta(2), 2\zeta(3), \dots)\right] \frac{(-y_1)^n}{n!}.$$

Let us build the noncommutative generating series of  $\zeta_{\sqcup}(w)$  and let us take the constant part of the two members of  $H(N) \xrightarrow[N \rightarrow \infty]{} \text{Const}(N) \pi_Y Z$ , we have

**Theorem 5.** Let  $Z_{\sqcup} = \sum_{w \in Y^*} \zeta_{\sqcup}(w)$   $w$  be the noncommutative generating series of the constants  $\zeta_{\sqcup}(w)$ . Then

$$Z_{\sqcup} = \left[ 1 + \sum_{n \geq 1} \left( \sum_{k=1}^n b_{n,k}(\gamma, \zeta(2), 2\zeta(3), \dots) \right) \frac{(-y_1)^n}{n!} \right] \pi_Y Z.$$

Identifying coefficients of  $y_1^k w$  in each member leads to

**Corollary 4.** For all  $w \in Y^* \setminus y_1 Y^*$  and  $k \geq 0$ , we have

$$\zeta_{\sqcup}(y_1^k w) = \sum_{i=1}^k \frac{\zeta_{\sqcup}(y_1^{k-i} w)}{i!} \left[ (-1)^i \sum_{j=1}^i b_{i,j}(\gamma, \zeta(2), 2\zeta(3), \dots) \right].$$

In consequence,

**Theorem 6** ([?]). For  $w \in Y^* \setminus y_1 Y^*$ ,  $k \geq 0$ , the constant  $\zeta_{\sqcup}(y_1^k w)$  associated to  $\zeta(y_1^k w)$  is a polynomial of degree  $k$  in  $\gamma$  and with coefficients in  $\mathcal{Z}$ . Moreover, for  $l = 0, \dots, k$ , the coefficient of  $\gamma^l$  is of weight  $|w| + k - l$ .

**Corollary 5.** For  $s > 1$ , the constant  $\zeta_{\sqcup}(1, s)$  associated to  $\zeta(1, s)$  is linear in  $\gamma$  and with coefficients in  $\mathbb{Q}[\zeta(2), \zeta(2i+1)]_{0 < i \leq (s-1)/2}$ .

**Example 3.**  $\gamma = \frac{\zeta_{\sqcup}(1, 2) + 2\zeta(3)}{\zeta(2)} = \frac{\zeta_{\sqcup}(1, 3) + \frac{1}{2}\zeta(2)^2}{\zeta(3)} = \frac{\zeta_{\sqcup}(1, 4) + 3\zeta(5) - \zeta(2)\zeta(3)}{\frac{2}{5}\zeta(2)^2}$ .

In other words, if we give to  $\gamma$  the weight<sup>1</sup> 1, then the constant  $\zeta_{\sqcup}(y_1^k w)$  associated to  $\zeta(y_1^k w)$  would be an homogeneous polynomial of weight  $|w| + k$ .

**Example 4.**

$$\begin{aligned} \zeta_{\sqcup}(1, 7) &= \zeta(7)\gamma + \zeta(3)\zeta(5) - \frac{54}{175}\zeta(2)^4, \\ \zeta_{\sqcup}(1, 1, 6) &= \frac{4}{35}\zeta(2)^3\gamma^2 + \left( \zeta(2)\zeta(5) + \frac{2}{5}\zeta(3)\zeta(2)^2 - 4\zeta(7) \right) \gamma \\ &\quad + \zeta(6, 2) + \frac{19}{35}\zeta(2)^4 + \frac{1}{2}\zeta(2)\zeta(3)^2 - 4\zeta(3)\zeta(5), \\ \zeta_{\sqcup}(1, 1, 1, 5) &= \frac{3}{4}\zeta(6, 2) - \frac{14}{3}\zeta(3)\zeta(5) + \frac{3}{4}\zeta(2)\zeta(3)^2 + \frac{809}{1400}\zeta(2)^4 \\ &\quad - \left( 2\zeta(7) - \frac{3}{2}\zeta(2)\zeta(5) + \frac{1}{10}\zeta(3)\zeta(2)^2 \right) \gamma + \left( \frac{1}{4}\zeta(3)^2 - \frac{1}{5}\zeta(2)^3 \right) \gamma^2 + \frac{1}{6}\zeta(5)\gamma^3. \end{aligned}$$

<sup>1</sup> In the theory of periods,  $\gamma$  is conjectured to be non-period and so would be transcendent.

### 2.2.3 Commutative generating series of polyzêtas

In Proposition ??, we explained how to extract the constant part of a noncommutative generating series of polyzêtas. Let us have a look now at following commutative generating series and corresponding to Ecalle's *Zig*-moulds [?]:

$$\mathcal{Z}(t_1, \dots, t_r) = \sum_{s_1, \dots, s_r > 0} \zeta(s_1, \dots, s_r) t_1^{s_1-1} \dots t_r^{s_r-1} = \sum_{n_1 > \dots > n_r > 0} \frac{1}{(n_1 - t_1) \dots (n_r - t_r)}. \quad (33)$$

These commutative generating series can be encoded by series  $\{S_j\}_{j=1, \dots, r}$  (or their projection over the alphabet  $Y$ ) of the form [?]

$$S_j = (t_j x_0)^* x_1 \dots (t_r x_0)^* x_1 = \sum_{s_j, \dots, s_r > 0} x_0^{s_j-1} x_1 \dots x_0^{s_r-1} x_1 t_j^{s_j-1} \dots t_r^{s_r-1}, \quad (34)$$

$$\pi_Y S_j = \sum_{s_j, \dots, s_r > 0} y_{s_j} \dots y_{s_r} t_j^{s_j-1} \dots t_r^{s_r-1} = \left( \sum_{s_j \geq 1} y_{s_j} t_j^{s_j-1} \right) \dots \left( \sum_{s_r \geq 1} y_{s_r} t_r^{s_r-1} \right). \quad (35)$$

Moreover, let  $S_{r+1} = 1$ . The series  $\mathcal{Z}(t_1, \dots, t_r)$  contain divergent terms of which we are looking for the constant part. We start from the following identity [?] due to convolution theorem [?]

$$S_1 = x_1^r + \sum_{j=1}^r t_j \sum_{i=0}^{j-1} x_1^i \sqcup x_0 [(-x_1)^{j-1-i} \sqcup S_j], \quad (36)$$

$$\Rightarrow \pi_Y S_1 = y_1^r + \sum_{j=1}^r t_j \left( \sum_{s_j \geq 2} y_1^{j-1} y_{s_j} t_j^{s_j-1} \right) \pi_Y S_{j+1}. \quad (37)$$

In consequence,

**Proposition 6.**

$$\zeta_{\sqcup}(S_1) = \sum_{j=1}^r (-1)^{j-1} t_j \zeta[x_0(x_1^{j-1} \sqcup S_j)].$$

$$\zeta_{\sqcup}(\pi_Y S_1) = (-1)^r \sum_{\substack{s_1, \dots, s_r > 0 \\ s_1 + \dots + s_r = r}} \frac{(-\gamma)^{s_1}}{s_1! \dots s_r!} \prod_{j=2}^r \left( -\frac{\zeta(j)}{j} \right)^{s_j} + \sum_{j=1}^r \sum_{s_j \geq 2} t_j t_j^{s_j-1} \zeta_{\sqcup}(y_1^{j-1} y_{s_j} \pi_Y S_{j+1}).$$

In the following part, iterated integrals associated with words  $w \in X^*$ , along the path  $0 \rightsquigarrow z$  and over differential forms  $\omega_0, \omega_1$ , will be denoted, as in [?], by  $\alpha_0^z(w)$ .

**Example 5.** Since  $(tx_0)^*x_1 = x_1 + tx_0(tx_0)^*x_1$ ,

$$\alpha_0^z[(tx_0)^*x_1] = \alpha_0^z(x_1) + t \int_0^z \left(\frac{z}{s}\right)^t \text{Li}_1(s) \frac{ds}{s} = \alpha_0^z(x_1) + t \sum_{n \geq 1} \frac{z^n}{n(n-t)}.$$

we get

$$\begin{aligned} \sum_{s \geq 1} \zeta_{\sqcup}(s)t^{s-1} &= \sum_{s \geq 2} \zeta(s)t^{s-1} = \sum_{n \geq 1} \left[ \frac{1}{n-t} - \frac{1}{n} \right], \\ \sum_{s \geq 1} \zeta_{\sqcup}(s)t^{s-1} &= \gamma + \sum_{s \geq 2} \zeta(s)t^{s-1} = \gamma + \sum_{n \geq 1} \left[ \frac{1}{n-t} - \frac{1}{n} \right]. \end{aligned}$$

**Example 6.** Identity (??) gives

$$S = x_1^2 + t_1 x_0(t_1 x_0)^*x_1(t_2 x_0)^*x_1 + t_2 x_0(-x_1 \sqcup (t_2 x_0)^*x_1) + t_2 x_1 \sqcup x_0(t_2 x_0)^*x_1.$$

In the second member of the previous expression,

- we have  $\zeta_{\sqcup}(x_1^2) = 0$ ,
- the first noncommutative rational series encodes, by convolution theorem [?], the following convergent integral

$$\begin{aligned} \alpha_0^1[(t_1 x_0)^*x_0 x_1(t_2 x_0)^*x_1] &= \int_0^1 \left(\frac{1}{s}\right)^{t_1} \frac{ds}{s} \int_0^s \frac{dr}{1-r} \sum_{n \geq 1} \frac{r^n}{n-t_2} \\ &= \sum_{n, m \geq 1} \frac{1}{(n+m)(n+m-t_1)(n-t_2)} \\ &= \sum_{n_1 > n_2 \geq 1} \frac{1}{n_1(n_1-t_1)(n_2-t_2)}, \end{aligned}$$

- the following rational series encodes

$$\begin{aligned} \alpha_0^1[x_0(-x_1 \sqcup (t_2 x_0)^*x_1)] &= - \int_0^1 \frac{ds}{s} \sum_{m \geq 1} \frac{s^m}{m} \sum_{n \geq 1} \frac{s^n}{n-t_2} \\ &= - \sum_{n, m \geq 1} \frac{1}{(n-t_2)(n+m)n} \\ &= - \sum_{n_1 > n_2 \geq 1} \frac{1}{n_1(n_1-n_2)(n_2-t_2)}, \end{aligned}$$

- the last noncommutative rational series corresponds to (but it has to be shuffled with  $x_1$  for which  $\zeta_{\sqcup}(x_1) = 0$ )

$$\alpha_0^1[x_0(t_2x_0)^*x_1] = \int_0^1 \frac{ds}{s} \sum_{n \geq 1} \frac{s^n}{n - t_2} = \sum_{n \geq 1} \frac{1}{n(n - t_2)}.$$

Thus,

$$\sum_{s,r \geq 1} \zeta_{\sqcup}(s,r)t_1^{s-1}t_2^{r-1} = \sum_{n_1 > n_2 \geq 1} \frac{t_1}{n_1(n_1 - t_1)(n_2 - t_2)} - \sum_{n_1 > n_2 \geq 1} \frac{t_2}{n_1(n_1 - n_2)(n_2 - t_2)}.$$

Projecting  $S$  over alphabet  $Y$ , we get successively, since  $y_1y_s = y_1 \sqcup y_s - y_{s+1} - y_sy_1$

$$\begin{aligned} \pi_Y S &= y_1^2 + \sum_{s,r \geq 2} y_sy_rt_1^{s-1}t_2^{r-1} + \sum_{s \geq 2} y_1y_st_2^{s-1} + \sum_{s \geq 2} y_sy_1t_1^{s-1} \\ &= y_1^2 + \sum_{s,r \geq 2} y_sy_rt_1^{s-1}t_2^{r-1} + y_1 \sqcup \sum_{s \geq 2} y_st_2^{s-1} - \frac{1}{t_2} \left[ y_1 + \sum_{s \geq 2} y_st_2^{s-1} \right] + \sum_{s \geq 2} y_sy_1[t_1^{s-1} - t_2^{s-1}] \\ &= y_1^2 - y_1t_2^{-1} + (y_1 - t_2^{-1}) \sqcup \sum_{s \geq 2} y_st_2^{s-1} + \sum_{s,r \geq 2} y_sy_rt_1^{s-1}t_2^{r-1} + \sum_{s \geq 2} y_sy_1[t_1^{s-1} - t_2^{s-1}]. \end{aligned}$$

In consequence,

$$\begin{aligned} \sum_{s,r \geq 1} \zeta_{\sqcup}(s,r)t_1^{s-1}t_2^{r-1} &= \frac{\gamma^2 - \zeta(2)}{2} - \frac{\gamma}{t_2} + (\gamma - t_2^{-1}) \sum_{n \geq 1} \left[ \frac{1}{n - t_2} - \frac{1}{n} \right] \\ &+ \sum_{n_1 > n_2 \geq 1} \left[ \frac{1}{n_1 - t_1} - \frac{1}{n_1} \right] \left[ \frac{1}{n_2 - t_2} - \frac{1}{n_2} \right] + \sum_{s \geq 2} \zeta(s, 1)[t_1^{s-1} - t_2^{s-1}]. \end{aligned}$$

The last sum can be encoded by  $x_0(t_1x_0)^*x_1^2 - x_0(t_2x_0)^*x_1^2$  and can be obtained from

$$\alpha_0^1[x_0(t_ix_0)^*x_1^2] = \int_0^1 \frac{ds}{s} \sum_{n_1 > n_2 \geq 1} \frac{s^{n_1}}{(n_1 - t_i)n_2} = \sum_{n_1 > n_2 \geq 1} \frac{1}{n_1(n_1 - t_i)n_2}.$$

## Appendix : application to polysystems [?]

Let  $q_1, \dots, q_n$  be commutative indeterminates over  $\mathbb{C}$ . We denote  $Q = \{q_1, \dots, q_n\}$ . The algebra of formal power series (resp. polynomials) over  $Q$  with coefficients in  $\mathbb{C}$  is denoted by  $\mathbb{C}[[Q]]$  (resp.  $\mathbb{C}[Q]$ ). An element of  $\mathbb{C}[[Q]]$  is an infinite sum  $f = \sum_{i_1, \dots, i_n \geq 0} f_{i_1, \dots, i_n} q_1^{i_1} \cdots q_n^{i_n}$ .

**Definition 3.** Let  $f \in \mathbb{C}[[Q]]$ . We set

$$\begin{aligned} E(f) &= \{\rho \in \mathbb{R}_+^n : \exists C_{f\rho} \in \mathbb{R}_+ \text{ such that for all } i_1, \dots, i_n \geq 0, |f_{i_1, \dots, i_n}| \rho_1^{i_1} \dots \rho_n^{i_n} \leq C_{f\rho}\} \\ \check{E}(f) &: \text{interior of } E(f) \text{ in } \mathbb{R}^n. \\ \text{CV}(f) &= \text{convergence domain of } f = \{q \in \mathbb{C}^n : (|q_1|, \dots, |q_n|) \in \check{E}(f)\}. \end{aligned}$$

The power series  $f$  is to be said convergent if  $\text{CV}(f) \neq \emptyset$ . Let  $\mathcal{U}$  be an open of  $\mathbb{C}^n$  and let  $q \in \mathbb{C}^n$ . The power series  $f$  is to be said convergent on  $q$  (resp. over  $\mathcal{U}$ ) if  $q \in \text{CV}(f)$  (resp.  $\mathcal{U} \subset \text{CV}(f)$ ). We set  $\mathbb{C}^{\text{cv}}[[Q]] = \{f \in \mathbb{C}[[Q]] : \text{CV}(f) \neq \emptyset\}$ . Let  $q \in \text{CV}(f)$ . There exist some constants  $C_{f\rho}, \rho$  and  $\check{\rho}$  such that  $|q_1| < \check{\rho} < \rho, \dots, |q_n| < \check{\rho} < \rho$  and  $|f_{i_1, \dots, i_n}| \rho^{i_1 + \dots + i_n} \leq C_{f\rho}$ , for  $i_1, \dots, i_n \geq 0$ . The convergence module of  $f$  at  $q$  is  $(C_{f\rho}, \rho, \check{\rho})$ .

Recall that  $D_1^{j_1} \dots D_n^{j_n} f$  is the partial derivation of order  $j_1, \dots, j_n \geq 0$  of  $f$  and is given by

$$\frac{D_1^{j_1} \dots D_n^{j_n} f}{j_1! \dots j_n!} = \sum_{i_1 \geq j_1, \dots, i_n \geq j_n} f_{i_1, \dots, i_n} \prod_{l=1}^n \binom{i_l}{j_l} q_n^{i_l - j_l}.$$

**Definition 4.** The polysystem  $\{A_i\}_{i=0, \dots, m}$  is defined by the Lie derivations  $A_i = \sum_{j=1}^n A_i^j D_j$ , where  $A_i^j \in \mathbb{C}^{\text{cv}}[[Q]]$ . It is linear if there exist  $\{M_i\}_{i=0, \dots, m} \in \mathcal{M}_{n,n}(\mathbb{C})$  s.t.

$$A_i = (q_1 \quad \dots \quad q_n) M_i \begin{pmatrix} D_1 \\ \vdots \\ D_n \end{pmatrix}.$$

Let  $f \in \mathbb{C}^{\text{cv}}[[Q]]$  and let  $\{A_i\}_{i=0,1}$  be a polysystem. Let  $(\rho, \check{\rho}, C_f)$  and let  $(\rho, \check{\rho}, C_i)$ , for  $i = 0, 1$ , be convergence modules of  $f$  and  $\{A_i^j\}_{j=1, \dots, n}$  respectively at  $q \in \text{CV}(f) \cap \bigcap_{i=0,1,j=1, \dots, n} \text{CV}(A_i^j)$ . We denote by  $(A_i f)|_q$  the evaluation at  $q$  of  $A_i f$ . Let us consider the system

$$y(z) = f(q(z)) \quad \text{where} \quad dq(z) = A_0(q)\omega_0(z) + A_1(q)\omega_1(z). \quad (38)$$

The generating series,  $\sigma f|_{q(z_0)}$ , associated to ?? and the Chen series  $S_{z_0 \rightsquigarrow z}$  are given by

$$\sigma f|_{q(z_0)} = \sum_{w \in X^*} A_w f|_{q(z_0)} w \quad \text{and} \quad S_{z_0 \rightsquigarrow z} = \sum_{w \in X^*} \alpha_{z_0}^z(w) w, \quad (39)$$

where  $A_w = \text{Id}$ ,  $\alpha_{z_0}^z(w) = 1$  if  $w = \epsilon$  and  $A_w = A_v A_i$ ,  $\alpha_{z_0}^z(w) = \int_{z_0 \rightsquigarrow z} \alpha_{z_0}^t(v) \omega_i(t)$  if  $w = vx_i$ . Since  $S_{z_0 \rightsquigarrow z}, L(z)L^{-1}(z_0)$  satisfy (??) taking the same value at  $z_0$  then  $S_{z_0 \rightsquigarrow z} = L(z)L^{-1}(z_0)$ . Hence, the asymptotic behaviour of  $L$  in (??) gives [?]

$$S_{\epsilon \rightsquigarrow 1-\epsilon} \sim e^{-x_1 \log \epsilon} Z e^{-x_0 \log \epsilon} \quad \text{for } \epsilon \rightarrow 0^+, \quad (40)$$

and the output  $y$  is given by  $y(z) = \langle \sigma f|_{q(z_0)} || S_{z_0 \rightsquigarrow z} \rangle = \sum_{w \in X^*} A_w f|_{q(z_0)} \alpha_{z_0}^z(w)$  [?].

Let  $\eta = q(z_0)$  and suppose that  $f(q) = \lambda q$  with  $\lambda \in \mathcal{M}_{1,n}(\mathbb{C})$ . If  $\{A_i\}_{i=0,1}$  is linear, then, by Definition ??, let  $M_i = \mu(x_i)$ , for  $i = 0, 1$ . Thus,  $\sigma f|_{q(z_0)} = \sum_{w \in X^*} [\lambda \mu(w) \eta] w$  is a rational power series of representation  $(\lambda, \mu, \eta)$  and it is a generating series of the differential system of rank  $n$ , or equivalently of the linear differential equation of order  $n$  with singularities in  $\{0, 1, \infty\}$ .

**Example 7** (hypergeometric equation).

$$z(1-z)\ddot{y}(z) + [t_2 - (t_0 + t_1 + 1)z]\dot{y}(z) - t_0 t_1 y(z) = 0.$$

Let  $q_1(z) = y(z)$  and  $q_2(z) = z(1-z)\dot{y}(z)$ . One has

$$\begin{pmatrix} dq_1 \\ dq_2 \end{pmatrix} = \left[ \begin{pmatrix} 0 & 0 \\ -t_0 t_1 & -t_2 \end{pmatrix} \omega_0 - \begin{pmatrix} 0 & 1 \\ 0 & t_2 - t_0 - t_1 \end{pmatrix} \omega_1 \right] \begin{pmatrix} q_1 \\ q_2 \end{pmatrix}.$$

Here  $y = (1 \ 0) \begin{pmatrix} q_1 \\ q_2 \end{pmatrix}$ ,  $M_0 = - \begin{pmatrix} 0 & 0 \\ t_0 t_1 & t_2 \end{pmatrix}$ ,  $M_1 = \begin{pmatrix} 0 & 1 \\ 0 & t_2 - t_0 - t_1 \end{pmatrix}$  and  $\eta = \begin{pmatrix} q_1(z_0) \\ q_2(z_0) \end{pmatrix}$ .

Thus, the solution of these equations can be obtained by examining the linear representation of generating series. The Drinfel'd equation allows to study the asymptotic behaviour, the functional equations and to compute the monodromy groups, the Galois differential groups.

**ACKNOWLEDGMENTS.** We acknowledge the influence of the Cartier's lectures at the GdT *Polylogarithmes et Polyzêtas*. We greatly appreciated many fruitful discussions with Enjalbert, Huttner, Jacob, Petitot and Waldschmidt.

## References

- [1] Hoang Ngoc Minh, Jacob, G., Oussous, N.E., and Petitot, M.: De l'algèbre des  $\zeta$  de Riemann multivariées à l'algèbre des  $\zeta$  de Hurwitz multivariées, *journal électronique du Séminaire Lotharingien de Combinatoire*, 44, 2001.
- [2] Hardy, G.H.: *Divergent series*, Clarendon, 1949.
- [3] Reutenauer, C.: *Free Lie Algebras*, London Math. Soc. Monographs, New Series-7, Oxford Science Publications, 1993.
- [4] Berstel, J. and Reutenauer, C.: *Rational series and their languages*, Springer-Verlag, 1988.
- [5] Hoffman, M.: The algebra of multiple harmonic series, *Journal of Algebra*, 1997.
- [6] Hoang Ngoc Minh: Finite polyzêtas, Poly-Bernoulli numbers, identities of polyzêtas and noncommutative rational power series, *Proceedings of 4<sup>th</sup> International Conference on Words*, 232-250, 2003.
- [7] Hoang Ngoc Minh, Petitot, M. and Van der Hoeven, J.: Polylogarithms and Shuffle Algebra, *Proceedings of FPSAC'98*, 1998.
- [8] Drinfel'd: Quasi-Hopf Algebras, *Leningrad Math. J.*, 1, 1419-1457, 1990.

- [9] Drinfel'd: On quasitriangular quasi-hopf algebra and a group closely connected with  $\text{gal}(\bar{q}/q)$ , *Leningrad Math. J.*, 4, 829-860, 1991.
- [10] Hoang Ngoc Minh, Petitot, M., and Van der Hoeven, J.: L'algèbre des polylogarithmes par les séries génératrices, *Proceedings of FPSAC'99*, 1999.
- [11] Lê, T.Q.T. and Murakami, J.: Kontsevich's integral for Kauffman polynomial, *Nagoya Math.*, 39-65, 1996.
- [12] Hoang Ngoc Minh: Shuffle algebra and differential Galois group of colored polylogarithms, *Nuclear Physics B*, 2004.
- [13] Costermans, C., Enjalbert, J.Y. and Hoang Ngoc Minh: Algorithmic and combinatoric aspects of multiple harmonic sums, *Discrete Mathematics & Theoretical Computer Science Proceedings*, 2005.
- [14] Ecalle, J.: *La libre génération des multizêta et leur décomposition canonico-explicite en irréductibles*, séminaire IHP, 1999.
- [15] Hoang Ngoc Minh: Summations of Polylogarithms via Evaluation Transform, *Mathematics and Computers in Simulations*, 42, 707-728, 1996.